



2005 ANNUAL UPDATE REPORT:

**BREEDING BIRD MONITORING IN
GREAT LAKES NATIONAL FORESTS:
1991-2005**

REPORT TO: CHEQUAMEGON/NICOLET,
CHIPPEWA AND SUPERIOR NATIONAL FORESTS

BY: JIM LIND, NICK DANZ, JOANN M.
HANOWSKI, AND GERALD J. NIEMI

NATURAL RESOURCES RESEARCH INSTITUTE
5013 MILLER TRUNK HIGHWAY
DULUTH, MN 55811

THIS IS NRRI TECHNICAL REPORT:
NRRI/TR-2006/06



SUMMARY

- A total of 132, 134, and 168 stands (1,257 survey points) were surveyed for breeding birds in the Chequamegon, Chippewa, and Superior National Forests (NF), respectively in 2005. Annual surveys have been conducted since 1991 in the Chippewa and Superior NF, and since 1992 in the Chequamegon NF.
- Trends in relative abundance were calculated for 69 bird species, including 57 species in the Chequamegon NF, 55 in the Chippewa NF, and 46 in the Superior NF. Thirty-nine species were also tested for a pooled trend by combining data from the three national forests.
- A total of 158 species/national forest trends were calculated in 2005 (not including pooled trends), 61 (39%) of which were significant ($P \leq 0.05$). Twenty species increased significantly ($P \leq 0.05$) in at least one national forest and 23 species decreased. Eight species had significant increasing pooled trends and 13 had decreasing trends. Of the 154 species/national forest trends calculated in 2004, 15 (10%) changed in 2005.
- The percentage of increasing species on each national forest ranged from 7% in the Chequamegon NF, to 22% in the Superior NF. The percentage of decreasing species ranged from 18% in the Chippewa NF, to 26% in the Chequamegon NF.
- To make additional comparisons of regional forest bird population trends, we also analyzed data from the Nicolet National Forest bird monitoring program in northeastern Wisconsin. Of the 71 species tested for trends between 1991 and 2004, four species increased significantly and 29 species declined.
- The short-distance migrant guild showed highly significant declines on all national forests. Long-distance migrants declined in the Chequamegon and Superior NF and increased in the Chippewa NF. Permanent residents increased on the Chippewa and Superior NF and were stable on the Chequamegon NF.
- The ground nesting guild declined on all national forests, while shrub/sub-canopy nesters increased on all national forests. The canopy and cavity nesting guilds showed stable trends, except for a decrease in canopy nesters in the Chequamegon NF and an increase in cavity nesters in the Superior NF.
- The deciduous and mixed forest bird guilds declined on the Chequamegon and Superior NFs and the pooled NFs. The lowland coniferous forest bird guild declined on the Chippewa NF and the pooled NFs. The early-successional bird guild increased on the Chippewa NF and the upland coniferous bird guild increased on the Superior NF.
- Evidence from recent regional studies have demonstrated greater nest predation rates on ground nests near forest/clearcut edges, as well as a significant increase in the creation of forest edges in recent years. Increasing amounts of forest edge and nest predation may be having negative effects on declining ground-nesters such as the Winter Wren, Veery, Hermit Thrush, Ovenbird, and White-throated Sparrow.

- Of the 1274 survey sites on the three national forests, 15.2% have been at least partially harvested since the beginning of monitoring, which is about 1% a year. This harvest rate is comparable to the documented 4.8% change from mature forest to early-successional types on federally managed forest lands in northeastern Minnesota between 1990 and 1995 (i.e. ~1% annual change). Thus, it appears that management activities on our sample sites are representative of the national forests as a whole, and that the trends we are documenting are probably occurring across the regional landscape.
- Many of the declining trends that we have detected have been consistent across the years and are not likely due to annual variation. One of the main goals of this monitoring program is to identify potential declines of forest bird species. This is especially true for species of conservation concern such as the Eastern Wood-Pewee, Winter Wren, Hermit Thrush, Ovenbird, and White-throated Sparrow. The declines observed over the past years for common species such as the Ovenbird and White-throated Sparrow are a continuing concern and special management consideration should be given to these species.

INTRODUCTION

The national forests of the western Great Lakes have among the richest diversity of breeding bird species in North America (Green 1995, Rich et al. 2004). An increased appreciation of this diversity, along with concerns about potential declines of some species, has led to a strong interest in monitoring forest bird populations in the region. The relatively heavily forested landscapes of northern Minnesota and Wisconsin are considered to be population “sources” for many forest bird species and may be supplementing population “sinks” in the agricultural landscapes of the lower Midwest (Robinson et al. 1995, Temple and Flaspohler 1998), highlighting the importance of monitoring trends in forest bird populations in the upper Midwest.

Agencies such as the USDA Forest Service have a need for population trend data at the scale of an individual national forest, to identify when and where population changes are occurring and identify potential conservation problems. Large-scale population monitoring programs such as the U.S. Geological Survey’s Breeding Bird Survey (BBS) provide important information on trends at a continental scale, however, limited coverage in some areas can make it difficult to use BBS data to characterize population trends at smaller geographic scales (Peterjohn et al. 1995). Continental trends also have the potential to mask regional population trends (Holmes and Sherry 1988), thus there is a need for regional monitoring programs that can provide more localized information (Green 1995, Howe et al. 1997).

In response to the need for regional population data, a long-term forest breeding bird monitoring program was established in 1991 on the Chippewa and Superior National Forests, and in 1992 on the Chequamegon National Forest and the St. Croix region of east-central Minnesota. The Forest Service is mandated to monitor certain management indicator species (Manley 1993), and our monitoring program expands beyond indicator species to include all forest songbird species that we can adequately sample. Currently, 435 stands (1,271 points) within the three national forests are surveyed once during each

breeding season (June 1 to July 10). From 1995 to 2001 we surveyed an additional 211 points in southeast Minnesota, however, counts were discontinued due to a lack of funding. See Lind et al. (2001b) for 1995-2001 results from southeast Minnesota. Surveys in the St. Croix region of east-central Minnesota were also discontinued after 2003 due to lack of funding, with 1992-2003 results available in our 2003 annual report (Lind et al. 2003). Results from the Nicolet National Forest bird monitoring program in northeastern Wisconsin will also be presented, to make additional comparisons of regional forest bird population trends.

The primary objective of this report is to update U.S. Forest Service personnel on results of the forest bird monitoring program. We focus on relative abundance trends of individual species, as well as assemblages of species, over the 14 to 15 year time frame of the monitoring. Our intent is to summarize the most important results and to provide detailed information in appendix form for those who need more specific results. This report, as well as annual update reports from 1998 to 2004, can be found on the internet at: <http://www.nrri.umn.edu/mnbirds/reports.htm>. Other objectives, including bird/habitat and bird/landscape relationships, development of management recommendations for birds, and development and monitoring of the forest plan, were met through Minnesota's Forest Bird Diversity Initiative (Niemi et al. 2003). Additional information on these objectives will be available as time and monetary resources become available.

DESIGN AND METHODS

Sample Design

The monitoring program was designed to provide an accurate estimate of population change for forest bird species on three national forests in northern Minnesota and Wisconsin (Figure 1). The spatial extent of each national forest is large, on the order of tens of thousands of hectares, and each area includes a mosaic of forest stand types. We distributed sampling locations across the forest mosaic in a stratified random manner. A list of forest stands was created for each study area, and stands with the same stand type according to dominant tree species and stocking density were grouped into strata. Stands were ≥ 16 ha (40 acres) and were identified from the individual national forest inventories. For each national forest, a number of stands were selected from each stratum so that the final proportion of stands of each stand type was equal to the proportion of forested land area of each stand type (Hanowski and Niemi 1995). Our sample of stands is therefore representative of the forest cover in each national forest. A total of 133, 135, and 169 stands were established in the Chequamegon, Chippewa, and Superior National Forests, respectively.

Stands were large enough to accommodate three sampling points a minimum of 220 meters apart. Changes to forest cover through natural and anthropogenic disturbance have occurred on sampling locations since the beginning of the study and may have caused concomitant changes in bird populations. Because sampling locations are permanently marked, we are able to incorporate such changes into our descriptions of bird population patterns through time.

Sampling

Point count sampling used in our program follow national and regional standards (Ralph et al. 1993, 1995, Howe et al. 1997). Ten-minute point counts were conducted at each point between June and early July (Reynolds et al. 1980). Point counts are appropriate for determining the relative abundance of most singing passerine species, but are inadequate for waterfowl, grouse, woodpeckers, and most raptors. In addition, because our surveys are conducted during the summer months, we may underestimate the relative abundance of early-nesting species (e.g. permanent residents that begin breeding in April, such as woodpeckers and chickadees).




Figure 1. Locations of forest breeding bird point counts in northern Minnesota and Wisconsin (1991-2005).

Point counts were conducted by trained observers (see observer training section below) from approximately 0.5 hour before to 4 hours after sunrise on days with little wind (< 15 km/hr) and little or no precipitation. All birds heard or seen from the point were recorded with estimates of their distance from that point. From 1991 to 1994, all birds heard or seen within 100 m of the point were recorded. From 1995-2005, we included all birds heard or seen from the point, regardless of distance, so that our results could be compared with other monitoring programs in this region (see Howe et al. 1997). The number of individuals observed for each species can be summed for 3, 5, and 10-minute

periods so that regional comparisons are possible with data gathered using 3 or 5-minute point counts.

We attempted to have each observer sample a similar number of stands of each forest cover type. This was done to minimize bias due to observer differences in sampling different forest cover types. Weather data (cloud cover, temperature, and wind speed) and time of day were recorded before each count.

Observer training

Prior to the field season, tapes of 120+ bird songs were provided as a learning tool, and all observers were required to pass an identification test of 75 bird songs made by Cornell University's Laboratory of Ornithology. A standard for number of correct responses was established by giving the test to observers who were trained in identifying birds by sound, and who had four to five years of field experience. This was done to identify songs on the tape that were not good representations of songs heard in northern Minnesota and Wisconsin. Based on results of trained observers, the standard for passing was set at 85% correct responses. Songs on the tape were grouped by habitat (e.g., upland deciduous, lowland coniferous) to simulate field cues that would aid in song identification.

Observer field training was conducted during the last week of May in the Superior National Forest. Observers conducted simultaneous practice counts at several points used in the monitoring program. Data were compiled for each observer, and species lists and numbers of individuals recorded on the count by each observer were compared to that of experienced observers. Deviations from the average or species missed were noted on the field sheets and returned. In addition to field training and testing, all observers were required to have a hearing test to ensure that their hearing was within normal ranges, as established by audiologists, for all frequencies (125 to 8000 hertz).

Analysis

The pattern of population change through time can be viewed in two distinct ways: 1) as *population trajectory*, the path of a population through time, including its ups and downs, and 2) as *population trend*, the overall pattern of increase or decrease over the course of the study, presented as a positive or negative number. We built statistical models of species relative abundance as a function of time to describe these features of bird populations.

Relative abundance

For each species, yearly relative abundance was calculated using birds detected within 100 m of each point. Relative abundance for species from the three national forests was calculated by summing the number of individuals of each species across two points per stand. In order to avoid double-counting of individuals, data from the two farthest separated points within a stand were summed and analyzed.

We used a set of criteria to ensure that our analyses provided reliable population information. Stands were included in the analysis only if they had been sampled in at least six years. Data were included for a species if it was observed on a minimum of five stands per study area and in at least three years on each stand. For species that were observed on a minimum of five stands in each of the three national forests, we pooled all stands and carried out an additional (three national forest combined) analysis. Although this pooled

analysis does not include lands in non-federal ownerships, it should give an indication of population trends at a larger scale than the individual national forest.

Population trajectory

Population trajectory can be thought of simply as the size of a population across time. Because we do not record every individual bird present in our study areas, we cannot know true population size. Instead, we must rely on our sample design to give an estimate of population size in each year. Central to our analytical process is how we scaled up bird abundance recorded at the stand level to an annual index of population size for the study areas. We used a non-parametric route regression procedure similar to that described by James et al. (1996), in which observed abundances on each stand are smoothed and then combined to give a region-wide index of population size.

We used locally-weighted (LOESS) regression to smooth the time series of species relative abundance for each stand. In LOESS-regression, fitted values (points along the curve) for years are calculated by giving a small amount of weight to neighboring years, for example, a year with high raw abundance for a species would tend to bring up the fitted values for the year before and the year after. We then computed the arithmetic mean and 95% confidence intervals using the fitted values from the within-stand regressions for each species in each year. The mean fitted value represents the annual index of population size. By plotting the mean fitted values and confidence intervals in a time series, we get a graphic depiction of the population trajectory (Appendix A). With every new year of sampling, we can expect the modeled abundance of a species in a given year to vary slightly from previous years' results, due to the way fitted abundance values are calculated in the LOESS-regression.

Population trend

Population trend can be thought of as a statement of the direction and magnitude of population change a given time period (Link and Sauer 1997). Because a significant trend implies a unidirectional change, linear methods can be used to detect trend without asserting that the population trajectory is linear (Urquhart and Kincaid 1999). To assess trend, we modeled the relationship between the annual index of population size for a study area (described in *Population Trajectory* above) and time using simple linear regression. We used the slope coefficient to characterize direction and magnitude of the trend. To facilitate comparison, slopes were converted to units of % annual change by dividing annual population indexes by the predicted value of the index at the midpoint of the survey period prior to regressing the index against time (Bart et al. 2003). We assessed the significance of the regressions using a bootstrap procedure (Manly 1990) in which trends were computed for 500 bootstrap resamples of the stands used to calculate the annual population index. For each bootstrap resample, trend was calculated using the same steps as for the original trend. For each original trend, an exact p-value was calculated as the percentile at which zero occurred in the distribution of 500 bootstrapped slopes. For example, $p = 0.01$ would be equivalent to 99% of bootstrapped slopes being greater than zero, which would give us a high degree of confidence that the true population slope was different from zero.

Guild Analyses

We examined trends for three types of guilds: migration strategy, nesting substrate, and vegetation-type preference (Appendix C). Guild analyses followed similar procedures as the individual species analyses, except that each species was assigned a guild category and all species within that category (e.g. long-distance migrants) were combined and analyzed as a group. All non-flyover individuals of all species within the 100 m radius were included, regardless of whether the species met the inclusion criteria described above for individual species. Guild categories were taken from Erlich et al. (1988) and Freemark and Collins (1992), with modifications based on personal experience and data from the region.

Note that some species use different migration strategies, nesting substrates, and vegetation types in different portions of their geographic range. Guild analyses also can be complicated by a lack of agreement on how to categorize guilds, and there will always be species that use multiple guilds. Species guilds are not mutually exclusive and the species pool in a migration guild, for example, can be very similar to the species pool in a nesting guild (Sauer et al. 1996). Directional trends in abundant species can strongly affect all the guilds that those species are categorized in. Given these limitations, we still feel it is important to look for underlying similarities among groups of increasing and decreasing species.

Methodological Considerations for Nicolet NF

The field protocols of the NRRI and Nicolet National Forest monitoring programs are similar enough to enable us, with a few minor considerations, to use Nicolet NF bird data in our trend analysis programs. Full methodological details of the Nicolet NF program can be found in Howe and Roberts (2005). Briefly, the Nicolet NF has carried out 10-minute unlimited-radius point counts on over 500 roadside and off-road (habitat-based) points since 1987. Individual points are surveyed in alternate years, with points in the southern portion of the forest surveyed in odd years since 1987 and northern points surveyed in even years since 1988.

To make NRRI and Nicolet NF trend results comparable, we limited our analysis of Nicolet NF data to habitat-based points (i.e. we excluded roadside points) between the years of 1991-2004. We used the exact same trend analysis programs with Nicolet NF data as described above for NRRI data. Howe and Roberts (2005) reported trends separately for the northern and southern portions of the forest. Alternatively, we computed forest-wide trends on the Nicolet by 1) carrying out LOESS regression to develop a smoothed time series of abundance for each species on each point, 2) obtaining fitted (predicted) values for years in which points were not surveyed (i.e. odd years for northern sites and even years for southern sites) and 3) calculating an annual index of abundance for each species by averaging the fitted values in each year using all points. The overall forest-wide trend was computed by regressing the annual index of abundance on time (year) as described above. We were limited to using 2004 as the most recent year for the Nicolet NF because northern sites were not surveyed in 2005 and the smoothing regression could not be used to extrapolate beyond the survey period; otherwise, the annual index for 2005 would be incorrectly computed from southern points only.

Trends were calculated for 71 forest breeding species present on a minimum of 5 Nicolet NF points. Species not well sampled by point count methodology (e.g. waterfowl, shorebirds, raptors, etc.) were excluded from analyses. Additionally, we computed trends for the total number of individuals in nesting, migration, and habitat guilds.

Range of Natural Variability

Bird population sizes in the Chippewa and Superior National Forests have been assessed in relation to their range of natural variability (RNV; Hanowski and Danz 2003). In an effort to place the trends from these two national forests into context with their historic and current populations, we have presented the RNV values for each of the species tested in the Chippewa and Superior NF. The RNV calculations are based on estimated historic forest conditions and the habitat affinities of each species, and represent a potential benchmark for evaluating our trends. An underlying assumption of the RNV concept is that a species is considered sustainable if it is currently present on the landscape at the same levels which it occurred historically.

RESULTS AND DISCUSSION

Over the course of 15 field seasons we have detected over 278,000 individual birds of 173 species on more than 18,500 ten-minute point counts in the three national forests (Figure 2). In 2005, we sampled 132 stands in the Chequamegon NF, 134 stands in the Chippewa NF, and 168 in the Superior NF.

Sixty-nine species were tested for trends in at least one national forest, including 57 in the Chequamegon NF, 55 in the Chippewa NF, and 46 in the Superior NF (Table 1). Additionally, 39 species were tested for a “pooled” (three national forests combined) trend. As monitoring has proceeded through the years, new species have met our criteria for inclusion in trend analyses on each national forest. Since these criteria were first applied in the 2000 analyses, the number of species tested has risen from 50, 47, 40, and 36, respectively. See Appendix A for graphs of individual species trajectories and Appendix B for test statistics and sample sizes used in the trend analyses.

Overview of Population Trends

A total of 158 species/national forest trends were calculated (not including pooled trends), 61 (39%) of which were significant ($P \leq 0.05$). Twenty species increased in at least one national forest, including four (Red-eyed Vireo, Black-capped Chickadee, Cedar Waxwing, and American Redstart) that increased in multiple national forests (Tables 2 and 3). Twenty-three species decreased in at least one national forest, including nine (Eastern Wood-Pewee, Great Crested Flycatcher, Winter Wren, Veery, Hermit Thrush, Black-throated Green Warbler, Ovenbird, Scarlet Tanager, and White-throated Sparrow) that decreased in multiple national forests. Of the 154 species/national forest trends calculated in 2004 (Lind et al. 2004), 15 (10%) changed in 2005 (Table 4).

Many of the species we monitor exhibit large annual fluctuations in abundance, a phenomenon which has been documented on several other long-term studies (Virkkala 1991, Blake et al. 1994, Weslowski and Tomialojc 1997, Holmes and Sherry 2001). Long-term monitoring studies are important for differentiating between these short-term fluctuations and actual long-term trends. In previous years' results, we often saw species

with opposite trends in different study areas (e.g. five species in 2000 results; Lind et al. 2001a). After 15 years of sampling, there now appears to be “core” groups of consistently increasing species (e.g. Red-eyed Vireo, Black-capped Chickadee, American Redstart) and decreasing species (e.g. Eastern Wood-Pewee, Winter Wren, Veery, Hermit Thrush, Ovenbird, and White-throated Sparrow).

Chequamegon National Forest

Of the 57 species tested for trends in the Chequamegon NF, four species (7%) increased significantly and 15 (26%) have decreased (Figure 3). The Yellow Warbler and American Redstart have the greatest rates of annual increase (>10%). No species are showing new increasing trends this year, and one formerly increasing species, the Red-eyed Vireo, now has a non-significant trend (0.74% annual increase; $P = 0.052$).

The Eastern Wood-Pewee, Winter Wren, Veery, and Hermit Thrush are well-represented on the Chequamegon NF, but have some of the greatest declines (6-9% annually; Appendix B). The Red-winged Blackbird and Evening Grosbeak have the two greatest rates of decrease, but they are tested on just five and six stands, respectively, and their trends may be more susceptible to site-specific influences than other species. Both of these species are, however, showing substantial declines in other parts of their ranges (Sauer 2004). The Great Crested Flycatcher, Blue-headed Vireo, Mourning Warbler, and Scarlet Tanager have new declines in 2005, and the Brown Creeper and Golden-crowned Kinglet were declining in 2004, but are no longer showing significant declines (Table 4).

Chippewa National Forest

Of the 55 species tested in the Chippewa NF, 11 species (20%) increased significantly and ten (18%) decreased (Figure 3). The Cedar Waxwing has the highest annual rate of increase (11%), but its population trajectory (Appendix A) shows an increase since the mid-1990's with an increase in the variance about the mean. This is probably due to encountering flocks of this gregarious species during some point counts and none in others. The Black-capped Chickadee, Chestnut-sided Warbler, and American Redstart are well-represented species on the forest, with 4-5% annual increases. The Red-eyed Vireo has one of the lowest rates of increase among the significantly increasing species, but because of its wide distribution, the increase is probably occurring over a large portion of the forest. It has also had a substantial increase since 1998 (Appendix A). The Canada Warbler and American Goldfinch had new increasing trends on the Chippewa NF this year, and the Gray Catbird is no longer increasing significantly (Table 4).

The greatest rate of annual decrease in the Chippewa NF is that of the Connecticut Warbler (13%). Although it is sampled on only 14 stands, the species has declined consistently since 1991 and the stands it occurs on are spread across most of the forest. Well-represented species that are showing annual rates of decline of 4% or more include the Eastern Wood-Pewee, Great Crested Flycatcher, Winter Wren, Hermit Thrush, and Song Sparrow. The Ovenbird is declining at 3%/year, but its trend may be especially important given its widespread distribution. No species have new decreasing trends on the Chippewa NF in 2005, and the Brown-headed Cowbird is no longer showing a significant decline (Table 4).

Six species are moving toward their historic population levels (RNV) on the Chippewa NF. Five species are below their RNV but have increasing trends (Black-

capped Chickadee, American Robin, Cedar Waxwing, Black-and-white Warbler, and Canada Warbler), and one species, the Song Sparrow, is above its RNV but declining (Table 7). Conversely, six species are moving away from their historic population levels. These species are below their RNV and decreasing (Winter Wren, Hermit Thrush, Nashville Warbler, Ovenbird, Connecticut Warbler, White-throated Sparrow). Of the 11 species within their RNV, five are increasing and two are decreasing.

Superior National Forest

Of the 46 species tested in the Superior NF, ten species (22%) are increasing and 11 (24%) are decreasing (Figure 3). The Hairy Woodpecker has the highest rate of annual increase (14%) of any species in the Superior NF, but it is tested on just five stands and its trend may be more susceptible to site-specific influences than other species. The Black-throated Blue Warbler and Cedar Waxwing have the next highest rates of increase (10-11% annually) on the 11 and nine stands they are each tested on, respectively. The Black-capped Chickadee, Golden-crowned Kinglet, American Redstart, and Northern Parula are widespread species that are increasing at 4-9% annually. The Hairy Woodpecker, Golden-crowned Kinglet, and Nashville Warbler have new increasing trends this year, and the Swamp Sparrow was increasing in 2004 but now has a stable trend (Table 4).

The Tennessee Warbler has the greatest rate of decrease (17% annually) on the Superior NF, however, it was just abundant enough to be tested for trends during its peak in the mid-1990's, and has had only 14 detections since 1999. The Eastern Wood-Pewee, Rose-breasted Grosbeak, Veery, Winter Wren, and White-throated Sparrow are declining at 3-10% annually, and are widely distributed on the forest. No species have new significant declines this year on the Superior NF, and all species that were declining in 2004 still are in 2005 (Table 4.)

Seven species are moving toward their historic population levels (RNV) on the Superior NF. Seven species are below their RNV but have increasing trends (including Northern Parula, Black-capped Chickadee and American Redstart), and no species are above their RNV (Table 7). Seven species are below their RNV and decreasing (including White-throated Sparrow, Winter Wren, and Scarlet Tanager). Of the 14 species within their RNV, two are increasing and four are decreasing.

Pooled National Forests

Of the 39 species tested for a pooled national forests trend, eight species (21%) increased significantly and 13 (33%) decreased (Figure 3). The Hairy Woodpecker increased at 5% annually although its trend was not statistically significant. The American Redstart and Black-capped Chickadee increased at about 4% annually. The Chestnut-sided Warbler and Red-eyed Vireo had lower rates of increase (2.6% and 1.6%, respectively), but both are widespread on all of the national forests and their lower rates still represent substantial increases. The Golden-crowned Kinglet and Magnolia Warbler each had new increases this year (this is the first year a pooled trends has been calculated for Magnolia Warbler), and the Blue Jay and Blackburnian Warbler are no longer showing a significant increase (Table 4).

The Eastern Wood-Pewee had the greatest annual rate of decrease (6.5%), which along with the Winter Wren, Hermit Thrush, Veery, Scarlet Tanager, Song Sparrow and

Common Yellowthroat, has had a consistent downward trajectory since the early 1990's (Appendix A). The White-throated Sparrow has declined 4% annually, but its trajectory has remained essentially level since 1997. The Yellow-rumped Warbler has a new decline this year and the Brown Creeper and Nashville Warbler are no longer declining (Table 4).

Management Activities on Study Areas

Of the 1274 survey sites on the three national forests, 15.2% have been at least partially harvested since the beginning of monitoring, which is about 1% a year (Table 8). A small number of our monitoring points have also had prescribed burns since the start of monitoring, but this is usually done after harvest. This harvest rate is comparable to the 4.8% change from mature forest to early-successional types on federally managed forest lands in northeastern Minnesota between 1990 and 1995 (i.e. ~1% annual change; Wolter and White 2002). Thus, it appears that management activities on our sample sites are representative of the national forests as a whole, and that the trends we are documenting are probably occurring across the regional landscape.

Guild Analyses

Short-distance migrants (species that winter mainly north of Mexico) showed highly significant declines ($P \leq 0.01$) in each national forest (Table 5). The most abundant short-distance migrants in our analyses include White-throated Sparrow, American Robin, Hermit Thrush, and Yellow-rumped Warbler. Long-distance migrants (species that winter mainly south of the U.S./Mexico border) showed mixed results across our study areas, with declines in the Chequamegon and Superior NF's and an increase in the Chippewa NF. Abundant long-distance migrants included Ovenbird, Red-eyed Vireo, Nashville Warbler, and Chestnut-sided Warbler. Permanent residents increased on all study areas except the Chequamegon NF, where they were stable. Black-capped Chickadee, Blue Jay, and Red-breasted and White-breasted nuthatches are the most abundant permanent residents.

Ground nesting birds showed highly significant declines in all study areas, while shrub/sub-canopy nesters increased in all study areas (Table 5). Abundant ground-nesters include Ovenbird, Nashville Warbler, Veery, and White-throated Sparrow. The most common shrub and subcanopy-nesting species include Red-eyed Vireo, Chestnut-sided Warbler, and American Redstart. Canopy and cavity nesters showed stable trends, except for a decrease in canopy nesters in the Chequamegon NF and an increase in cavity nesters in the Superior NF. Most primary cavity excavators (e.g. woodpeckers) have had stable trends, while many secondary excavators (e.g. chickadees and nuthatches) have had increasing trends. An exception is the Great Crested Flycatcher (a secondary excavator) which is declining in the Chequamegon and Chippewa NF.

The deciduous and mixed forest bird guilds declined on the Chequamegon and Superior NFs and the pooled NFs. The lowland coniferous forest bird guild declined on the Chippewa NF and the pooled NFs. The early-successional bird guild increased on the Chippewa NF and the upland coniferous bird guild increased on the Superior NF.

Nicolet NF trend comparisons

Of the 71 species tested for trends on the Nicolet NF between 1991 and 2004, four species increased significantly and 29 species declined (Tables 1 & 6). Three of the four increasing Nicolet NF species were also increasing on at least one of the NRR study areas

(Red-eyed Vireo, Nashville Warbler, and American Redstart). Twenty-five of the 29 declining species were also tested for a trend on the NRRI monitoring program, and of these, ten were declining on at least one of the NRRI study areas. These include species such as the Eastern Wood-Pewee, Winter Wren, Veery, and White-throated Sparrow, which have had widespread declines documented on NRRI study areas. Conversely, five of the declining Nicolet NF species were increasing in at least one NRRI study area, including Black-capped Chickadee, White-breasted Nuthatch, and Cedar Waxwing (Table 1). Most of the remaining declining Nicolet NF species have relatively small sample sizes on NRRI study areas (Table 6).

The disparity in results between the Nicolet NF and NRRI bird monitoring programs is difficult to interpret. There are some minor differences in methodology, such as sampling conducted by volunteers during a single weekend and the use of unlimited radius point counts on the Nicolet NF, but given the proximity to the Chequamegon NF (<50 miles away) the two monitoring programs should be sampling similar bird populations and landscapes. Although there are some similarities in trends from the two monitoring programs, the Nicolet NF has documented a greater proportion of declining species (41%) and a lower proportion of increasing species (5%) than any of the national forests in the NRRI program (Figure 3). However, the proportion of increasing species on the Chequamegon NF (the closest forest to the Nicolet NF) has been dropping in recent years to about 7% in 2005, with about 25% of the tested species declining. It will be interesting to see if future trends in forest bird populations become more or less similar among these two monitoring programs.

Conclusions

Most of the species with widespread increasing trends are either forest habitat generalists (e.g. Red-eyed Vireo, Black-capped Chickadee and Blue Jay) or early successional species (e.g. Cedar Waxwing, Chestnut-sided Warbler and American Redstart). Many of these increasing species are currently at or above their estimated RNV values. Recent increases in the amount of edge and early-successional habitat on the regional landscape (Wolter and White 2002) may be benefiting these species. The Black-capped Chickadee is a year-round resident that may also be responding to increased food availability from bird feeding activities, especially considering their increasing numbers on Minnesota Christmas Bird Counts in the past decade (National Audubon Society 2004).

The Blackburnian Warbler is a mature coniferous/mixed forest species that has also shown widespread increases. Population fluctuations in this species are often attributed to changes in spruce budworm (*Choristoneura fumiferana*) abundance. There was an outbreak in early 1990's with a decline since 1998 (Blackford 2001), that seems to correspond to the Blackburnian Warbler's trajectory (Appendix A). However, this is difficult to corroborate with other spruce budworm specialists (e.g. Tennessee, Bay-breasted and Cape May warblers) which are on the southern fringe of their ranges in our study areas.

Species with widespread declines on our study sites are mainly found in mature forest habitats, with the possible exception of Veery and White-throated Sparrow. While White-throated Sparrow abundance is often higher in clearcuts than in mature forests, reproductive rates have been shown to be up to three times greater in older forests (75-100

years) than in younger forests (Rangen et al. 2000). The Eastern Wood-Pewee, Winter Wren, Veery, and White-throated Sparrow have each shown significant declines on our surveys as well as USGS Breeding Bird Survey routes over much of their range (Sauer 2004). Increases in edge and early-successional habitats may be having negative effects on these species, although there are examples of increases in mature forest species on individual national forests (e.g., White-breasted Nuthatch Black-throated Blue Warbler, Northern Waterthrush).

The declines in ground nesters and increases in shrub nesters in our study seem to occur irrespective of migration strategy and habitat. It is possible that declines in ground-nesting populations are being influenced by recent changes in the landscapes of the Upper Midwest. Although the landscape surrounding the three national forests is primarily forested, average forest stand sizes and ages have changed in recent years. Wolter and White (2002) demonstrated a substantial decrease in patch size and interior forest area and a significant increase in edge density in early successional forest types in northeastern Minnesota between 1990 and 1995. Studies have shown that nesting success is reduced in landscapes with reduced patch sizes and high amounts of edge habitat, probably due to an increase in generalist nest predators (Robinson et al. 1995, Donovan et al. 1997). In the forested landscapes of the upper Midwest, recent studies have found higher predation rates on ground nests near forest/clearcut edges than in interior areas (Fenske-Crawford and Niemi 1997, Manolis et al. 2000, Flaspohler et al. 2001). Data from the Minnesota DNR winter track survey (Berg 2001) between 1991 and 2000 indicate a peak in track indices in 1995 for potential ground nest predators such as fisher (*Martes pennati*) and pine marten (*Martes martes*), which loosely follows the declines between 1994 and 1996 in many of the species we monitor. Nonetheless, the effects of nest predation on population trends in this study are unknown.

Many of the declining trends that we have detected have been consistent across the years and are not likely due to annual variation. Many of the declining trends that we have detected have been consistent across the years and are not likely due to annual variation. One of the main goals of this monitoring program is to identify potential declines of forest bird species. This is especially true for species of conservation concern such as the Eastern Wood-Pewee, Winter Wren, Hermit Thrush, Ovenbird, and White-throated Sparrow. The declines observed over the past years for common species such as the Ovenbird and White-throated Sparrow are a continuing concern and special management consideration should be given to these species. Several species are currently well below their estimated RNV values and they may not remain common if their declining trends continue.

Many of the declining species breed in mature forests, and many are ground-nesters. Some of these population declines may be linked to recent reductions in forest patch size and stand age on the landscape, especially in light of regional studies showing high nest predation on ground-nests near forest edges. Although the factors responsible for population declines are not definitively known, the prominence of declining ground-nesting species suggests that it would be prudent to curb further reductions in average forest patch sizes and age on the landscape. Several of these declining species have high PIF conservation values (e.g. Veery, Mourning Warbler, Eastern Wood-Pewee), and the extensive forests of northern Minnesota and Wisconsin represent excellent opportunities to provide “source” populations for many species.

LITERATURE CITED

- Bart, J., B. Collins, and R.I.G. Morrison. 2003. Estimating population trends with a linear model. *Condor* 105:367-372.
- Berg, B. 2001. Winter track survey summary, 2000. Forest Wildlife Populations and Research Group, Minnesota Department of Natural Resources. Grand Rapids, MN.
- Blackford, D. C. 2001. Spruce budworm project. Forest Insect and Disease Newsletter. July 2001. Minnesota Department of Natural Resources, Division of Forestry.
- Blake, J. G., J. M. Hanowski, G. J. Niemi, and P. T. Collins. 1994. Annual variation in bird populations of mixed conifer-northern hardwood forests. *Condor* 96:381-399.
- Donovan, T. M., P. J. Jones, E. M. Annand, and F. R. Thompson, III. 1997. Variation in local-scale edge effects: mechanisms and landscape context. *Ecology* 78:2064-2075.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. 785 pp. Simon and Schuster, Inc. New York, New York.
- Fenske-Crawford, T. J., and G. J. Niemi. 1997. Predation of artificial ground nests at two types of edges in a forest-dominated landscape. *Condor* 99:14-24.
- Flaspohler, D. J., S. A. Temple, and R. N. Rosenfield. 2001. Species specific edge effects on nest success and breeding bird density in a forested landscape. *Ecological Applications* 11:32-46.
- Freemark, K., and B. Collins. 1992. Landscape ecology of birds breeding in temperate forest fragments. Pages 443-454 in *Ecology and conservation of Neotropical migrant landbirds* (J.M. Hagan and D.W. Johnston, eds). Smithsonian Institution Press, Washington, D.C.
- Green, J. C. 1995. *Birds and forests: A management and conservation guide*. Minnesota Department of Natural Resources. St. Paul, MN. 182 pp.
- Hanowski, J. M., and N. Danz. 2003. Response of breeding birds to forest plan revision alternatives in the Chippewa and Superior National Forests. Report to USDA Forest Service – Superior and Chippewa National Forests (available at <http://www.nrri.umn.edu/mnbirds/reports.htm>).
- Hanowski, J. M., and G. J. Niemi. 1995. Experimental design considerations for establishing an off-road, habitat specific bird monitoring program using point counts. Pages 145-150 in *Monitoring bird populations by point counts*. General Technical Report PSW-GTR-149. Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture, Albany, CA.
- Holmes, R. T., and T. W. Sherry. 1988. Assessing population trends of New Hampshire forest birds: Local vs. regional trends. *Auk* 105:756-768.
- Holmes, R. T., and T. W. Sherry. 2001. Thirty-year bird population trends in an unfragmented temperate deciduous forest: importance of habitat change. *Auk* 118:589-609.

- Howe, R. W., G. J. Niemi, G. J. Lewis, and D. A. Welsh. 1997. A standard method for monitoring songbird populations in the Great Lakes region. *Passenger Pigeon* 59:182-194.
- James, F. C., C. E. McCulloch, and D. A. Wiedenfeld. 1996. New approaches to the analysis of population trends in land birds. *Ecology* 77:13-27.
- Lind, J., N. Danz, M. T. Jones, J. M. Hanowski, and G. J. Niemi. 2001a. Breeding bird monitoring in Great Lakes National Forests: 1991-2000. Natural Resources Research Institute Technical Report: NRRI/TR-2001/4.
- Lind, J., N. Danz, M. T. Jones, J. M. Hanowski, and G. J. Niemi. 2001b. Breeding bird monitoring in Great Lakes National Forests: 1991-2001. Natural Resources Research Institute Technical Report: NRRI/TR-2001/39. (available at <http://www.nrri.umn.edu/mnbirds/reports.htm>)
- Lind, J., N. Danz, M. T. Jones, J. M. Hanowski, and G. J. Niemi. 2002. Breeding bird monitoring in Great Lakes National Forests: 1991-2002. Natural Resources Research Institute Technical Report: NRRI/TR-2002/24. (available at <http://www.nrri.umn.edu/mnbirds/reports.htm>)
- Lind, J., N. Danz, J. M. Hanowski, and G. J. Niemi. 2004. Breeding bird monitoring in Great Lakes National Forests: 1991-2004. Natural Resources Research Institute Technical Report: NRRI/TR-2005/04. (available at <http://www.nrri.umn.edu/mnbirds/reports.htm>)
- Link, W. A., and J. R. Sauer. 1997. New approaches to the analysis of population trends in land birds: comment. *Ecology* 78:2632-2634.
- Lynch, J. F., and D. F. Whigham. 1984. Effects of forest fragmentation on breeding bird communities in Maryland, U.S.A. *Biological Conservation* 28:287-324.
- Manley, P. N. and Monitoring Task Group. 1993. Guidelines for monitoring populations of neotropical migratory birds on National Forest System lands. U.S. For. Serv. Wildlife and Fisheries. U.S. Govt. Printing Office 1993-720-803/80195.
- Manly, B.F.J. 1991. Randomization and Monte Carlo methods in biology. Chapman & Hall, London, UK.
- Manolis, J. C., D. E. Andersen, and F. J. Cuthbert. 2000. Patterns in clearcut edge and fragmentation effect studies in northern hardwood-conifer landscapes: retrospective power analysis and Minnesota results. *Wildlife Society Bulletin* 28:1088-1101.
- MathSoft, Inc. 1999. S-Plus User's Guide, Data Analysis Products Division, MathSoft, Seattle, WA. 634 pp.
- National Audubon Society. 2004. The Christmas Bird Count historical results [Online]. Available <http://www.audubon.org/bird/cbc> [accessed Jan. 2005].
- Niemi, G. J., J. M. Hanowski, N. Danz, J. Lind, M. Jones, and J. Sales. 2003. Minnesota's Forest Bird Diversity Initiative. Natural Resources Research Institute Technical Report: NRRI/TR-2003/11.

- Peterjohn, B. G., J. R. Sauer, and C. S. Robbins. 1995. Population trends from the North American Breeding Bird Survey. Pages 3-39 in Ecology and management of neotropical migratory birds (T. E. Martin and D. M. Finch, eds.). Oxford University Press, New York.
- Ralph, C. J., G. R. Geupel, P. Pyle, T. E. Martin, and D. F. DeSante. 1993. Handbook of field methods for monitoring landbirds. Gen. Tech. Rep. PSW-GTR-144. Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture, Albany, CA. 41 pp.
- Ralph, C. J., J. R. Sauer, and S. Droege (eds.). 1995. Monitoring bird populations by point counts. Gen. Tech. Rep. PSW-GTR-149. Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture, Albany, CA. 181 pp.
- Rangen, S.A., K.A. Hobson, and R.G. Clark. 2000. A comparison of density and reproductive indices of songbirds in young and old boreal forest. *Wildlife Society Bulletin* 28:110-118
- Reynolds, R. T., J. M. Scott, and R. A. Nussbaum. 1980. A variable circular-plot method for estimating bird numbers. *Condor* 82:309-313.
- Rich, T.D., C.J. Beardmore, H. Berlanga, P.J. Blancher, M.S. Bradstreet, G.S. Butcher, D.W. Demarest, E.H. Dunn, W.C. Hunter, E.E. Itigo-Elias, J.A. Kennedy, A.M. Martell, A.O. Panjabi, D.N. Pashley, K.V. Rosenberg, C.M. Rustay, J.S. Wendt, T.C. Will. 2004. Partners In Flight North American Landbird Conservation Plan. Cornell Lab of Ornithology, Ithaca, NY.
- Robinson, S. K., F. R. Thompson III, T. M. Donovan, D. R. Whitehead, and J. Faaborg. 1995. Regional forest fragmentation and the nesting success of migratory birds. *Science* 267:1987-1990.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2004. The North American Breeding Bird Survey, Results and Analysis 1966 - 2003. Version 2004.1, USGS Patuxent Wildlife Research Center, Laurel, MD. <http://www.mbr-pwrc.usgs.gov/bbs/bbs.html>
- Sauer, J. R., G. W. Pendleton, and B. G. Peterjohn. 1996. Evaluating causes of population change in North American insectivorous songbirds. *Conserv. Biology* 10:465-478.
- Temple, S. A., and D. J. Flaspohler. 1998. The edge of the cut: implications for wildlife populations. *Journal of Forestry* 96:22-26.
- Urquhart, N. S., and T. M. Kincaid. 1999. Designs for detecting trend from repeated surveys of ecological resources. *Journal of Agricultural, Biological, and Environmental Statistics* 4:404-414.
- Virkkala, R. 1991. Spatial and temporal variation in bird communities and populations in north-boreal coniferous forests: a multiscale approach. *Oikos* 62:59-66.
- Weslowski, T., and L. Tomialojc. 1997. Breeding bird dynamics in a primeval temperate forest: long-term trends in Bialowieza National Park (Poland). *Ecography* 20:432-453.
- Wolter, P. T., and M. A. White. 2002. Recent forest cover type transitions and landscape structural changes in northeast Minnesota. *Landscape Ecology* 17:133-155.

Table 1. Trends for three individual national forests (NF), combined NF's, and Nicolet NF based on linear regression (1991-2005). I = significantly increasing, D = significantly decreasing. * $P \leq 0.05$, ** $P \leq 0.01$. See Appendix A for species graphs and Appendix B for test statistics and sample sizes.

Species	Chequamegon NF	Chippewa NF	Superior NF	Combined Trend	Nicolet NF
Yellow-bellied Sapsucker	ns	ns	ns	ns	ns
Downy Woodpecker		ns			D*
Hairy Woodpecker	ns	ns	I**	ns	ns
Northern Flicker	ns		ns		D**
Olive-sided Flycatcher		ns			D**
Eastern Wood-Pewee	D**	D**	D**	D**	D**
Yellow-bellied Flycatcher	ns	ns	ns	I**	ns
Alder Flycatcher	ns	ns	ns	ns	ns
Least Flycatcher	ns	I*	ns	ns	D**
Eastern Phoebe					ns
Great Crested Flycatcher	D**	D**			D**
Eastern Kingbird	ns				D**
Yellow-throated Vireo	ns	ns			ns
Blue-headed Vireo	D*	ns	ns	ns	ns
Warbling Vireo					ns
Red-eyed Vireo	ns	I**	I*	I**	I**
Gray Jay		ns	ns		ns
Blue Jay	ns	ns	ns	ns	ns
Black-capped Chickadee	ns	I**	I**	I**	D**
Red-breasted Nuthatch	ns	ns	ns	ns	ns
White-breasted Nuthatch	ns	I**			D*
Brown Creeper	ns	ns	ns	ns	ns
House Wren	ns				ns
Winter Wren	D**	D**	D**	D**	D**
Sedge Wren					D*
Golden-crowned Kinglet	ns	ns	I**	I**	ns
Eastern Bluebird					D**
Ruby-crowned Kinglet			D*		
Veery	D**	ns	D**	D**	D**
Swainson's Thrush			ns		ns
Hermit Thrush	D**	D**	ns	D**	ns
Wood Thrush	ns				ns
American Robin	ns	I*	ns	ns	ns
Gray Catbird		ns			ns
Brown Thrasher	ns				D**
Cedar Waxwing		I**	I**		D**
Golden-winged Warbler	ns	ns	ns	ns	D*
Tennessee Warbler			D**		

Table 1 (continued)

Species	Chequamegon NF	Chippewa NF	Superior NF	Combined Trend	Nicolet NF
Nashville Warbler	ns	D*	I*	ns	I**
Northern Parula	ns	ns	I**	I**	ns
Yellow Warbler	I**	D*			D**
Chestnut-sided Warbler	ns	I**	ns	I**	ns
Magnolia Warbler	ns	ns	I**	I**	ns
Cape May Warbler					ns
Black-throated Blue Warbler			I**		ns
Yellow-rumped Warbler	D*	ns	ns	D**	ns
Black-throated Green Warbler	D**	ns	D**	D*	ns
Blackburnian Warbler	I**	ns	ns	ns	ns
Pine Warbler	ns	ns	ns	ns	ns
Palm Warbler		ns			
Black-and-white Warbler	ns	I**	ns	ns	D**
American Redstart	I**	I**	I**	I**	I**
Ovenbird	D**	D**	D**	D**	ns
Northern Waterthrush	I*	ns	D**	ns	ns
Connecticut Warbler		D**			
Mourning Warbler	D*	ns	ns	D**	D**
Common Yellowthroat	D**	ns	ns	D*	ns
Canada Warbler	ns	I**	ns	ns	ns
Scarlet Tanager	D**	ns	D**	D**	D**
Eastern Towhee	ns				D*
Chipping Sparrow	ns	ns	ns	ns	ns
Clay-colored Sparrow	ns				
Vesper Sparrow	ns				
Field Sparrow					D**
Song Sparrow	ns	D**	ns	D**	D**
Lincoln's Sparrow					I*
Swamp Sparrow	ns	ns	ns	ns	ns
White-throated Sparrow	D*	D**	D**	D**	D**
Rose-breasted Grosbeak	ns	ns	D**	D**	D**
Indigo Bunting	ns	ns			D**
Red-winged Blackbird	D*	ns			ns
Common Grackle					ns
Brewer's Blackbird	ns				
Brown-headed Cowbird	ns	ns			D**
Baltimore Oriole					D**
Purple Finch		ns			D*
American Goldfinch	ns	I*			ns
Evening Grosbeak	D**				ns

Table 2. Species trends ($P \leq 0.05$) for three national forests (1991-2005), based on simple linear regression. ** $P \leq 0.01$. Species graphs can be found in Appendix A.

Increasing Species

<u>Chequamegon NF</u>	<u>Chippewa NF</u>	<u>Superior NF</u>	<u>Pooled national forests</u>
** Yellow Warbler	Least Flycatcher	** Hairy Woodpecker	** Yellow-bellied Flycatcher
** Blackburnian Warbler	** Red-eyed Vireo	Red-eyed Vireo	** Red-eyed Vireo
** American Redstart	** Black-capped Chickadee	** Black-capped Chickadee	** Black-capped Chickadee
Northern Waterthrush	** White-breasted Nuthatch	** Golden-crowned Kinglet	** Golden-crowned Kinglet
	American Robin	** Cedar Waxwing	** Northern Parula
	** Cedar Waxwing	Nashville Warbler	** Chestnut-sided Warbler
	** Chestnut-sided Warbler	** Northern Parula	** Magnolia Warbler
	** Black-and-white Warbler	** Magnolia Warbler	** American Redstart
	** American Redstart	** Black-throated Blue Warbler	
	** Canada Warbler	** American Redstart	
	American Goldfinch		

Decreasing Species

<u>Chequamegon NF</u>	<u>Chippewa NF</u>	<u>Superior NF</u>	<u>Pooled national forests</u>
** Eastern Wood-Pewee	** Eastern Wood-Pewee	** Eastern Wood-Pewee	** Eastern Wood-Pewee
** Great Crested Flycatcher	** Great Crested Flycatcher	** Winter Wren	** Winter Wren
Blue-headed Vireo	** Winter Wren	Ruby-crowned Kinglet	** Veery
** Winter Wren	** Hermit Thrush	** Veery	** Hermit Thrush
** Veery	Nashville Warbler	** Tennessee Warbler	** Yellow-rumped Warbler
** Hermit Thrush	Yellow Warbler	** Black-throated Green Warbler	Black-throated Green Warbler
Yellow-rumped Warbler	** Ovenbird	** Ovenbird	** Ovenbird
** Black-throated Green Warbler	** Connecticut Warbler	** Northern Waterthrush	** Mourning Warbler
** Ovenbird	** Song Sparrow	** Scarlet Tanager	Common Yellowthroat
Mourning Warbler	** White-throated Sparrow	** White-throated Sparrow	** Scarlet Tanager
** Common Yellowthroat		** Rose-breasted Grosbeak	** Song Sparrow
** Scarlet Tanager			** White-throated Sparrow
White-throated Sparrow			** Rose-breasted Grosbeak
Red-winged Blackbird			
** Evening Grosbeak			

Table 3. Summary of species with increasing or decreasing trends ($P \leq 0.05$) on three national forests (1991-2005). Individual species graphs can be found in Appendix A.

Increased in one national forest	Increased in two national forests	Increased in three national forests
Hairy Woodpecker	Red-eyed Vireo	American Redstart
Least Flycatcher	Black-capped Chickadee	
White-breasted Nuthatch	Cedar Waxwing	
Golden-crowned Kinglet		
American Robin		
Nashville Warbler		
Northern Parula		
Yellow Warbler		
Chestnut-sided Warbler		
Magnolia Warbler		
Black-throated Blue Warbler		
Blackburnian Warbler		
Black-and-white Warbler		
Northern Waterthrush		
Canada Warbler		
American Goldfinch		
Decreased in one national forest	Decreased in two national forests	Decreased in three national forests
Blue-headed Vireo	Great Crested Flycatcher	Eastern Wood-Pewee
Ruby-crowned Kinglet	Veery	Winter Wren
Tennessee Warbler	Hermit Thrush	Ovenbird
Nashville Warbler	Black-throated Green Warbler	White-throated Sparrow
Yellow Warbler	Scarlet Tanager	
Yellow-rumped Warbler		
Northern Waterthrush		
Connecticut Warbler		
Mourning Warbler		
Common Yellowthroat		
Song Sparrow		
Rose-breasted Grosbeak		
Red-winged Blackbird		
Evening Grosbeak		

Table 4. Summary of changes in trends on three national forests between 2004 and 2005 analyses.

<u>Species with new significant trends ($P \leq 0.05$) in 2005</u>			
<i>Increasing</i>		<i>Decreasing</i>	
<u>Species</u>	<u>Study area</u>	<u>Species</u>	<u>Study area</u>
Canada Warbler	Chippewa NF	Great Crested Flycatcher	Chequamegon NF
American Goldfinch	Chippewa NF	Blue-headed Vireo	Chequamegon NF
Hairy Woodpecker	Superior NF	Mourning Warbler	Chequamegon NF
Nashville Warbler	Superior NF	Scarlet Tanager	Chequamegon NF
Golden-crowned Kinglet	Superior NF	Yellow-rumped Warbler	Pooled NF's
Golden-crowned Kinglet	Pooled NF's		
Magnolia Warbler	Pooled NF's		
<u>Species no longer showing significant trends ($P > 0.05$) in 2005</u>			
<i>Was increasing in 2004</i>		<i>Was decreasing in 2004</i>	
<u>Species</u>	<u>Study area</u>	<u>Species</u>	<u>Study area</u>
Red-eyed Vireo	Chequamegon NF	Brown Creeper	Chequamegon NF
Gray Catbird	Chippewa NF	Golden-crowned Kinglet	Chequamegon NF
Swamp Sparrow	Superior NF	Brown-headed Cowbird	Chippewa NF
Blue Jay	Pooled NF's	Brown Creeper	Pooled NF's
Blackburnian Warbler	Pooled NF's	Nashville Warbler	Pooled NF's

Table 5. Test statistics and sample sizes for guild trend analyses on three national forests (1991-2005). All species combined within each guild category and analyzed as a group, regardless of whether a species meets criteria for individual species analyses. Change = percent annual change. N = number of stands analyzed. See Appendix A for trend graphs.

Guild Category		Chequamegon NF				Chippewa NF				Superior NF				Pooled National Forests			
		change	P	R ²	N	change	P	R ²	N	change	P	R ²	N	change	P	R ²	N
Migration	Short-distance	-2.625	0.000	0.916	129	-1.271	0.004	0.597	126	-1.522	0.000	0.621	147	-1.751	0.000	0.915	402
	Long-distance	-0.857	0.000	0.874	129	0.537	0.008	0.162	126	-0.486	0.024	0.408	147	-0.361	0.004	0.180	402
	Permanent Resident	-1.005	0.152	0.746	121	1.719	0.008	0.749	125	1.900	0.000	0.922	145	1.101	0.016	0.643	391
Nesting	Ground	-2.605	0.000	0.947	129	-1.959	0.000	0.598	126	-1.752	0.000	0.724	147	-2.210	0.000	0.765	402
	Shrub/Sub-canopy	0.603	0.228	0.169	129	3.001	0.000	0.914	126	0.691	0.040	0.643	147	1.472	0.000	0.818	402
	Canopy	-1.494	0.000	0.703	124	-0.596	0.180	0.349	126	-0.136	0.632	0.105	147	-0.633	0.008	0.826	397
	Cavity	-0.864	0.156	0.799	125	0.902	0.120	0.397	125	2.694	0.000	0.955	144	1.030	0.016	0.817	394
Vegetation Preference	Coniferous forest	-0.422	0.524	0.068	107	0.856	0.120	0.545	110	1.283	0.000	0.723	145	0.656	0.056	0.466	362
	Lowland coniferous	-0.952	0.124	0.267	107	-1.502	0.000	0.669	99	0.219	0.572	0.094	147	-0.737	0.000	0.355	353
	Deciduous forest	-1.482	0.000	0.818	127	-0.032	0.952	0.000	126	-0.784	0.008	0.657	147	-0.673	0.000	0.576	400
	Early-succession	0.841	0.464	0.587	101	2.417	0.004	0.751	118	-0.954	0.256	0.158	145	0.127	0.752	0.003	364
	Mixed forest	-3.370	0.000	0.982	120	-0.608	0.256	0.237	123	-1.820	0.000	0.809	147	-1.768	0.000	0.796	390

Table 6. Test statistics and sample sizes for individual species and guild trends from the Nicolet National Forest (1991-2004). Change = percent annual change. N = number of stands analyzed. See Appendix A for trend graphs.

Species	change	<i>P</i>	R ²	N
Yellow-bellied Sapsucker	-0.124	0.908	0.020	93
Downy Woodpecker	-4.936	0.048	0.837	17
Hairy Woodpecker	-0.265	0.836	0.008	24
Northern Flicker	-5.047	0.004	0.854	53
Olive-sided Flycatcher	-10.461	0.000	0.998	13
Eastern Wood-Pewee	-3.593	0.008	0.943	115
Yellow-bellied Flycatcher	-1.983	0.468	0.293	29
Alder Flycatcher	-0.297	0.872	0.045	54
Least Flycatcher	-3.429	0.000	0.975	124
Eastern Phoebe	-5.001	0.504	0.922	17
Great Crested Flycatcher	-6.284	0.000	0.989	117
Eastern Kingbird	-4.941	0.004	0.992	39
Yellow-throated Vireo	-4.728	0.384	0.845	17
Blue-headed Vireo	-3.886	0.196	0.832	41
Warbling Vireo	-1.652	0.456	0.718	11
Red-eyed Vireo	0.473	0.008	0.196	307
Gray Jay	-9.083	0.084	0.747	10
Blue Jay	-0.530	0.668	0.328	254
Black-capped Chickadee	-1.431	0.000	0.782	214
Red-breasted Nuthatch	-0.638	0.220	0.055	108
White-breasted Nuthatch	-6.039	0.032	0.601	32
Brown Creeper	-3.485	0.420	0.680	57
House Wren	-2.591	0.780	0.878	18
Winter Wren	-2.503	0.008	0.903	124
Sedge Wren	-1.975	0.048	0.527	9
Golden-crowned Kinglet	-2.948	0.276	0.974	46
Eastern Bluebird	-12.288	0.000	0.991	18
Veery	-3.651	0.004	0.955	104
Swainson's Thrush	-5.529	0.124	0.470	6
Hermit Thrush	-0.908	0.244	0.856	245
Wood Thrush	-2.228	0.132	0.748	35
American Robin	-0.264	0.176	0.225	262
Gray Catbird	-6.396	0.308	0.655	15
Brown Thrasher	-9.947	0.004	0.721	10
Cedar Waxwing	-5.744	0.008	0.972	133
Golden-winged Warbler	-5.380	0.040	0.869	44
Nashville Warbler	2.001	0.000	0.705	197
Northern Parula	-2.688	0.116	0.889	65
Yellow Warbler	-8.010	0.000	0.963	36
Chestnut-sided Warbler	-3.381	0.220	0.923	161
Magnolia Warbler	0.030	0.924	0.000	15

Table 6 (continued)

Species	change	<i>P</i>	R ²	N
Cape May Warbler	4.145	0.320	0.972	7
Black-throated Blue Warbler	-6.003	0.064	0.935	28
Yellow-rumped Warbler	-1.945	0.836	0.719	107
Black-throated Green Warbler	-0.279	0.516	0.432	185
Blackburnian Warbler	-0.266	0.460	0.018	66
Pine Warbler	-2.595	0.852	0.571	23
Black-and-white Warbler	-3.313	0.000	0.979	113
American Redstart	1.243	0.008	0.108	28
Ovenbird	0.900	0.064	0.828	291
Northern Waterthrush	-2.165	0.380	0.329	15
Mourning Warbler	-4.646	0.000	0.911	142
Common Yellowthroat	-1.916	0.164	0.929	106
Canada Warbler	-9.091	0.192	0.969	18
Scarlet Tanager	-8.033	0.000	0.982	114
Eastern Towhee	-7.826	0.012	0.866	17
Chipping Sparrow	-2.484	0.440	0.789	78
Field Sparrow	-9.474	0.000	0.952	8
Song Sparrow	-4.217	0.000	0.987	114
Lincoln's Sparrow	0.942	0.048	0.132	12
Swamp Sparrow	-1.174	0.652	0.618	53
White-throated Sparrow	-3.260	0.000	0.971	177
Rose-breasted Grosbeak	-2.158	0.000	0.820	244
Indigo Bunting	-5.377	0.000	0.960	86
Red-winged Blackbird	-0.744	0.052	0.306	63
Common Grackle	7.128	0.532	0.866	24
Brown-headed Cowbird	-9.405	0.004	0.981	35
Baltimore Oriole	-6.680	0.004	0.869	30
Purple Finch	-9.008	0.012	0.987	28
American Goldfinch	-4.064	0.112	0.962	102
Evening Grosbeak	-4.148	0.916	0.271	29
Guild Trends	change	<i>P</i>	R ²	N
Short-distance migrant	-2.379	0.000	0.991	318
Long-distance migrant	-1.107	0.004	0.888	318
Permanent resident	-1.771	0.020	0.941	306
Ground nester	-1.006	0.236	0.871	317
Shrub/subcanopy nester	-1.764	0.016	0.936	318
Canopy nester	-1.597	0.000	0.909	317
Cavity nester	-3.244	0.000	0.938	301
Coniferous forest	-1.723	0.108	0.931	260
Lowland coniferous forest	0.205	0.004	0.032	230
Deciduous forest	-0.753	0.000	0.918	318
Early successional forest	-3.855	0.000	0.958	249
Mixed forest	-1.638	0.000	0.940	307

Table 7. Comparison of species trends (1991-2005) and % of the range of natural variability (RNV) for 2003 populations on the Chippewa and Superior National Forests (from Hanowski and Danz 2003). 100% of RNV indicates that a species was considered to be within its historic range of natural variability. I = significantly increasing, D = significantly decreasing. * $P \leq 0.05$, ** $P \leq 0.01$.

Species	Chippewa NF trend	% of RNV	Superior NF trend	% of RNV
Yellow-bellied Sapsucker	ns	100%	ns	100%
Downy Woodpecker	ns	122%		
Hairy Woodpecker	ns	71%	I**	
Olive-sided Flycatcher	ns	53%		
Eastern Wood-Pewee	D**	100%	D**	100%
Yellow-bellied Flycatcher	ns	60%	ns	100%
Least Flycatcher	I*	100%	ns	100%
Great Crested Flycatcher	D**	100%		
Yellow-throated Vireo	ns	110%		
Blue-headed Vireo	ns	39%	ns	66%
Red-eyed Vireo	I**	100%	I*	97%
Gray Jay	ns	69%	ns	69%
Blue Jay	ns	72%	ns	69%
Black-capped Chickadee	I**	95%	I**	70%
Red-breasted Nuthatch	ns	56%	ns	70%
White-breasted Nuthatch	I**	100%		
Brown Creeper	ns	91%	ns	77%
Winter Wren	D**	60%	D**	87%
Golden-crowned Kinglet	ns	80%	I**	82%
Ruby-crowned Kinglet			D*	100%
Veery	ns	100%	D**	100%
Swainson's Thrush			ns	100%
Hermit Thrush	D**	70%	ns	68%
American Robin	I*	62%	ns	99%
Gray Catbird	ns	244%		
Cedar Waxwing	I**	88%	I**	87%
Golden-winged Warbler	ns	112%	ns	
Tennessee Warbler			D**	100%
Nashville Warbler	D*	52%	I*	83%
Northern Parula	ns	44%	I**	65%
Chestnut-sided Warbler	I**	100%	ns	100%
Magnolia Warbler	ns	38%	I**	100%
Black-throated Blue Warbler			I**	100%
Yellow-rumped Warbler	ns	46%	ns	63%
Black-throated Green Warbler	ns	76%	D**	80%

Table 7 (continued)

Species	Chippewa NF trend	% of RNV	Superior NF trend	% of RNV
Blackburnian Warbler	ns	70%	ns	88%
Pine Warbler	ns	56%	ns	72%
Palm Warbler	ns	28%		
Black-and-white Warbler	I**	59%	ns	100%
American Redstart	I**	100%	I**	78%
Ovenbird	D**	88%	D**	99%
Northern Waterthrush	ns		D**	51%
Connecticut Warbler	D**	97%		
Mourning Warbler	ns	100%	ns	100%
Canada Warbler	I**	47%	ns	100%
Scarlet Tanager	ns	105%	D**	58%
Chipping Sparrow	ns	88%	ns	82%
Song Sparrow	D**	110%	ns	86%
White-throated Sparrow	D**	42%	D**	89%
Rose-breasted Grosbeak	ns	100%	D**	97%
Indigo Bunting	ns	160%		
Brown-headed Cowbird	ns	120%		
Purple Finch	ns	62%		

Table 8. Number of survey sites harvested in each national forest since the start of monitoring.

Study Area	Total # of sites	# clearcut	# partially or selectively cut*	% harvested
Chequamegon NF	390	15	31	11.8%
Chippewa NF	393	21	49	17.8%
Superior NF	491	42	36	15.9%

* Sites in the partially cut category can include anywhere from 10-90% of the 100 m radius count circle harvested.






Figure 2. Total number of individuals detected annually (1991-2005) in each national forest (inside 100 m radius), based on raw data before applying analysis criteria (e.g. includes flyovers, etc.). The number of sites sampled is presented below each year.




Figure 3. Summary of significant trends ($P \leq 0.05$) by national forest (1991-2005). Pooled trends include three national forests combined. See Table 1 for list of species trends by national forest.

Appendix A

Trends in relative abundance by study area for all species and guilds tested (1991-2005)

Please see the *Analysis* section in the body of the report for details about how the plots were constructed.